Gastroenterology Therapies Please fax completed referral form to 972-473-7563

PATIENT DEMOGRAPHIC INFORMATION

atient's name:		Date of birth:	Phone #:	
ddress:		_ City/State/Zip:		
llergies:		Weight:	Height:	
		IICAL INFORMATION		
iagnosis: (ICD 10 Code Requir				
1 Crohn's Disease (K50.00-K50.919), ICD 10 1 Other:		☐ Ulcerative Colitis (K51.00-K51.919), ICD 10		
Other:				
rior Failed Therapy:				
		Reason for D/C:		
		Reason for D/C:		
ed Falled:	Length of Treatment:		Reason for D/C:	
est Results: Please attach cop	y for all items checked.			
l Negative TB test l Other:				
- Othor:				
Please include con	v of insurance card, most recen	t H&P or MD note. m	edication list, and lab results.	
T Todoo morado oop	y or modranico cara, most rocon			
	THERAPY	ORDERED		
MEDICATION	DOSE	DIRECTIONS/DURATION		
Cimzia® (certolizumab pegol)	INITIAL: 400mg	☐ INITIAL: Inject 400mg	g SUBQ at Weeks 0, 2, and 4	
	MAINTENANCE:		ct 400mg SUBQ every 4 weeks x 1 year	
	□ 400mg	☐ MAINTENANCE: Inje	ect 200mg SUBQ every 2 weeks x 1 year	
	□ 200mg	E INITIAL L.C IV	W I . 0 0 I 0	
Entyvio® (vedolizumab)	300mg	☐ INITIAL: Infuse IV at	weeks u, z, and 6 ise IV every 8 weeks x 1 year	
Remicade® (infliximab)	□ 5mg/kg: mg	□ INITIAL: Infuse IV at		
	□ 10mg/kg: mg		ise IV every 8 weeks x 1 year	
	☐ Other:mg		ise IV every weeks x 1 year	
Renflexis® (infliximab-abda)	□ 5mg/kg: mg	☐ INITIAL: Infuse IV at	Weeks 0, 2, and 6	
	□ 10mg/kg:mg		ise IV every 8 weeks x 1 year	
	☐ Other:mg	☐ MAINTENANCE: Infu	ise IV every weeks x 1 year	
Inflectra® (infliximab-dyyb)	□ 5mg/kg: mg	☐ INITIAL: Infuse IV at	Weeks 0, 2, and 6	
	□ 10mg/kg :mg		ise IV every 8 weeks x 1 year	
	☐ Other:mg	☐ MAINTENANCE: Infu	se IV every weeks x 1 year	
Tysabri® (natalizumab)	300mg	☐ Infuse IV every 4 wee	ks x 1 year	
OTHER:				
OTHER:				
patient currently receiving the	erapy above from another facility?	□ NO □ YES If yes, Fac	ility Name:	
		CIAN INFORMATION		
hysician Name:				
	Ci			
Office Contact:	Ph	ione #:	Fax #:	

